



ELSEVIER

NO signals in the haze

Nitric oxide signalling in plant defence

Margit Leitner, Elodie Vandelle, Frank Gaupels, Diana Bellin and Massimo Delledonne

Nitric oxide (NO) is gaining increasing attention as a regulator of diverse (patho-)physiological processes in plants. Although this molecule has been described as playing a role in numerous conditions, its production, turnover and mode of action are poorly understood. Recent studies on NO production have tended to highlight the questions that still remain unanswered rather than telling us more about NO metabolism. But regarding NO signalling and functions, new findings have given an impression of the intricacy of NO-related signalling networks. Different targets of protein S-nitrosylation have been characterised and enzymatic routes controlling this posttranslational modification are emerging, along with their physiological implications. Evidence is also accumulating for protein tyrosine nitration and cGMP as important components of NO-related signal transduction.

Address

Università degli Studi di Verona, Dipartimento di Biotecnologie, Strada Le Grazie 15, 37134 Verona, Italy

Corresponding author: Delledonne, Massimo (massimo.delledonne@univr.it)

Current Opinion in Plant Biology 2009, 12:451–458

This review comes from a themed issue on
Biotic Interactions
Edited by Xinnian Dong and Regine Kahmann

Available online 14th July 2009

1369-5266/\$ – see front matter
© 2009 Elsevier Ltd. All rights reserved.

DOI [10.1016/j.pbi.2009.05.012](https://doi.org/10.1016/j.pbi.2009.05.012)

Introduction

Nitric oxide (NO) is a small gaseous radical with diverse signalling functions. In plants, NO was first found to play a crucial role in mediating defence reactions against bacterial pathogens [1] and is now well known to influence numerous physiological processes throughout the entire plant life cycle. To name a few, NO is involved in germination, leaf expansion, lateral root development, flowering, stomatal closure, cell death and defence against biotic and abiotic stresses [2*]. Whereas descriptions of NO-mediated processes are accumulating, the plant signalling pathways governed by NO are still largely unknown. NO-related signalling can be attributed to various NO derivatives, collectively referred to as reactive

nitrogen species (RNS). RNS comprise not only the NO radical (NO^{*}) and its nitroxyl (NO⁻) and nitrosonium (NO⁺) ions, but also peroxyxynitrite (ONOO⁻), S-nitrosothiols, higher oxides of nitrogen and dinitrosyl-iron complexes; in short, all NO derivatives that can effect NO-dependent modifications [3]. Hence, the term NO-related signalling is used here to summarise effects caused by all these RNS.

In principle, NO-related functions are subject to different levels of control. Production and turnover regulate NO bioavailability; once the NO level increases in the system, it can affect signalling either directly via protein modifications or indirectly via activation of second messengers. In animals, second messengers, such as cGMP, are well-characterised components of NO signal transduction, whereas studies of NO-dependent posttranslational modifications are more recent. However, in research conducted in plants, knowledge of the cGMP-dependent pathways is restricted to data gained using pharmacological approaches. By contrast, the impact of NO-dependent protein modifications, especially of S-nitrosylation, is the best studied mode of action in plants to date.

Among the diverse physiological processes affected by NO, available data predominantly explain signalling related to plant defence responses—which is the focus of this review.

Regulation of NO bioavailability

No simple answers to the question how NO is produced in plants

Three routes to yield NO have been described in plants: non-enzymatic conversion of nitrite to NO in the apoplast, nitrate reductase (NR)-dependent NO formation and NO synthase (NOS)-like activity, that is arginine-dependent NO formation. These pathways have been reviewed in detail [2*,4*]. In a nutshell: since the enzymatic source(s) of NO in plant stress responses remains elusive, unbiased genetic tools are still lacking for non-invasive manipulations of NO levels *in planta*. Despite numerous studies using pharmacological tools demonstrated NOS-like activity in plants, the identity of the enzymes involved remains unknown. The only postulated plant NOS (AtNOA1/RIF1) has recently been shown to have no NOS activity [5**]. Instead, it is a chloroplast-targeted GTPase essential for proper ribosome assembly [6*]. Mutation in this gene leads to reduced NO accumulation, probably because of its rapid

Table 1

Characterised mutants and transgenics with altered NO levels.						
Gene	Description	Mod.	Species	NO level	related phenotype	Ref.
<i>Noa1/Rif1</i>	Nitric oxide associated1/resistant to fosmidomycin 1; chloroplastic GTPase	m	<i>At</i>	–	ROS accumulation	[5**,6*]
		s	<i>At</i>	+	–	[5**,6*]
<i>Nia1 Nia2</i>	Nitrate reductase	m	<i>At</i>	–	Reduced amino acid and nitrite levels	[8]
<i>NiR</i>	Nitrite reductase	as	<i>Nt</i>	+	Nitrite accumulation	[12]
<i>Argah1</i> and <i>2</i>	Arginine amidohydrolase; arginase	m	<i>At</i>	+	L-Arginine accumulation	[9]
<i>Nox1/cue1</i>	NO overproducer 1/chlorophyll a/b binding protein underexpressed 1; PEP phosphate translocator	m	<i>At</i>	+	L-Arginine accumulation	[10]
<i>Rcd1</i>	Radical-induced cell death1; unknown function.	m	<i>At</i>	+	ROS accumulation	[13]
<i>Gsnor1/hot5</i>	GSNO reductase	m	<i>At</i>	+	Accumulation of S-nitroso species and nitrate	[11*,19,20]
		as	<i>At</i>	+		
		s	<i>At</i>	–	–	[11*,19]
<i>Vsr3</i>	Vacuolar sorting receptor	as	<i>At</i>	+	ROS accumulation	[14]
<i>Hb1</i>	Ns haemoglobin	as	<i>At, Ms</i>	+	–	[17]
		s	<i>At, Ms</i>	–	Nitrate accumulation	[17]
		s	<i>At</i>	–	Nitrate accumulation	[18]

as, antisense; *At*, *Arabidopsis thaliana*; GSNO, S-nitrosoglutathione; Mod., gene modification; *Ms*, *Medicago sativa*; m, mutant; *Nt*, *Nicotiana tabacum*; Ns, non-symbiotic; PEP, phosphoenolpyruvate; Ref., references; ROS, reactive oxygen species; s, sense.

reaction with the elevated amounts of ROS observed in the *Atmoa1* mutant [5**].


By contrast, NO can clearly be produced by NR activity [7]. However, NR deficiency also leads to impaired nitrogen assimilation, thus generally influencing primary and secondary metabolism. Indeed, *Arabidopsis nia1 nia2* double mutants display decreased levels of both nitrite and amino acids [8]. Hence, direct effects due to impaired NO biosynthesis are difficult to distinguish from those caused by metabolic alterations. In several other mutants, increased NO accumulation correlates with concentrations of putative substrates for NO biosynthesis (Table 1): (1) Arginase-deficient *Atargah 1-1* and *2-1* as well as *Atnox1/cue1*, defective in a phosphoenolpyruvate phosphate translocator, display elevated levels of L-arginine [9,10]; (2) *Atgsnor1-3/hot5-2*, defective in the S-nitrosoglutathione reductase, accumulates nitrate and (3) an antisense *NiR* (nitrite reductase) tobacco accumulates nitrite [11*,12]. Recently, the *Arabidopsis rcd1*, a mutant sensitive to ozone and *VSR3* (vacuolar sorting receptor 3) antisense transgenic were shown to overaccumulate NO and ROS, extending the list of genetically modified plants with altered NO and/or ROS homeostasis [13,14]. In summary, none of these NO-related mutants is exclusively affected in NO production. Their complex and pleiotropic phenotypes make genetic study of NO function difficult. The significance of using genetic tools for studying NO-related signalling *in vivo* is underlined by recent reports demonstrating the strict co-localisation of NO sources (i.e. NOSs) and S-nitrosylation in mammalian systems [15,16]. Therefore, the identification and

characterisation of NO-producing enzymes in plants, other than NR, remain equally challenging and mandatory tasks.

Different paths lead to RNS and their turnover

Relative to the elusive routes of NO production, mechanisms regulating bioavailable RNS are comparatively well characterised. As outlined above, different RNS contribute to NO-related signalling. Accordingly, a network of RNS transformation and turnover balances the bioavailability of these signal compounds. These pathways are illustrated here using three examples. NO can be metabolised to nitrate by non-symbiotic haemoglobins, like *Arabidopsis Hb1*, which acts as an NO dioxygenase using NADPH as an electron donor (Figure 1, [17]). However, the NO-reducing activity of *Arabidopsis Hb1* is rather low, although physiologically relevant under hypoxic stress [17]. NO also reacts with GSH to form S-nitrosylated glutathione (GSNO), which can release NO or function as a transnitrosylating agent; it is thus considered a reactive nitrogen species (RNS) and natural reservoir of NO. The enzyme controlling GSNO levels is GSNO reductase (GSNOR). It reduces GSNO to oxidised glutathione and NH₃. Though highly specific for GSNO, GSNOR seems to also influence levels of protein S-nitrosylation [19]. In *Arabidopsis*, mutation of GSNOR (*gsnor1-3/hot5-2*) causes increased basal levels of NO and S-nitroso species. The mutant plants not only have impaired resistance to biotic and abiotic stresses, but also show reduced growth, fertility and reproduction. This complex phenotype underlines the vital role of functional RNS homeostasis in diverse physiological processes

Figure 1



Routes of NO production and turnover. Three pathways of NO production are thought to exist in plants: NOS-like activity of (an) unidentified protein(s), NO production by nitrate reductase and non-enzymatic conversion of nitrite to NO under acidic conditions. Three main routes lead to NO turnover: NO scavenging by non-symbiotic haemoglobins, transformation of NO to GSNO and degradation by GSNOR and reaction of NO with superoxide to form peroxynitrite, which is detoxified by peroxiredoxins. Abbreviations: GSNO, S-nitrosoglutathione; GSNOR, S-nitrosoglutathione reductase; NOS, nitric oxide synthase; nsHb, non-symbiotic haemoglobin; Prx, peroxiredoxin; RNS, reactive nitrogen species.


[11^{*},20,21]. Finally, NO rapidly reacts with superoxide (O_2^-) to form peroxynitrite ($ONOO^-$), a potent oxidising and nitrating species. This non-enzymatic reaction is tightly controlled by the (enzymatic) formation of its precursors. $ONOO^-$ can, in turn, be detoxified by some peroxiredoxins. For example, Arabidopsis PrxII E possesses peroxynitrite reductase activity (Figure 1) [22^{**}] and therefore controls the bioavailability of $ONOO^-$.

Hence, a complex interplay between RNS becomes evident: by removing NO, other RNS can be formed, presumably with altered signalling functions ([23], cf. Figure 1 and below). Moreover, peroxiredoxin II E is inhibited by S-nitrosylation, causing increased $ONOO^-$ -mediated protein tyrosine nitration. This demonstrates that enzyme regulation by NO can facilitate accumulation of another RNS.

NO-associated protein modifications

RNS can directly react with diverse biomolecules [3]. Amongst them, proteins can be modified by RNS through reactions with different amino acids or prosthetic groups.

Figure 2



Characterised targets of protein S-nitrosylation in plants. S-nitrosylation under physiological conditions (basal NO tone, blue arrows) is assumed to inhibit MAT1 and pMC9 and to facilitate oligomer formation of NPR1. Induced NO production (i.e. defence-related NO burst; red arrows) leads to S-nitrosylation of GAPDH, SABP3 and PrxII E. GAPDH activity is blocked upon S-nitrosylation and the protein is thought to translocate to the nucleus. S-nitrosylation of SABP3 leads to the loss of the enzyme's SA-binding capacity as well as inhibition of its carbonic anhydrase activity. Also PrxII E activity is inhibited by S-nitrosylation, which leads to increased peroxynitrite levels and protein nitration. Under induced conditions, thioredoxins counteract S-nitrosylation of NPR1, leading to monomer release and translocation of this transcription co-factor to the nucleus. Dashed lines indicate hypothetical interactions, S-nitrosylation is symbolised by red bullets. Abbreviations: GAPDH, glyceraldehyde 3-phosphate dehydrogenase; JA, jasmonic acid; MAT1, methionine adenosyltransferase 1; NPR1, non-expressor of pathogenesis-related protein 1; pMC9, prometacaspase 9; PrxII E, peroxiredoxin II E; SABP3, salicylic acid binding protein 3; TRX, thioredoxin.

More specifically, the main NO-associated protein modifications in the biological context are the covalent modifications of cysteine (S-nitrosylation) and tyrosine (tyrosine 3-nitration) residues and NO binding to transition metals (metal nitrosylation). To date, the best characterised of these is cysteine S-nitrosylation.

Protein S-nitrosylation

S-nitrosylation refers to the covalent attachment of an NO moiety to the thiol side chain of cysteine [24]. Although proteomic approaches identified numerous candidates in plants [25,26], only a few confirmed functional modifications have been reported (Figure 2). Nevertheless, beginning characterisation of modified proteins now reveals the physiological significance of S-nitrosylation.

S-nitrosylation contributes to gene regulation

NO-dependent gene regulation strongly depends on the plant organ and the inducing stimulus, reinforcing the

need to consider downstream regulatory networks [27]. Changes in gene expression have been intensively studied using different NO donors (reviewed by [28]), though the regulatory basis leading to these changes remains unknown. Common regulatory elements in differentially expressed genes could be identified using bioinformatics tools [29]. These theoretical considerations have recently been substantiated by experimental data that documented the control of NO over transcription factors [30,31**]. In particular, finding NPR1, a crucial component of disease resistance and signal cross-talk, to be regulated via *S*-nitrosylation added an important clue to understanding NO's signalling functions [31**]. The subcellular localisation of NPR1, and thus its transcription co-factor activity, is controlled by *S*-nitrosylation. Intermolecular disulphide bridges sustain NPR1 oligomers, which are retained in the cytosol. Intriguingly, *S*-nitrosylation facilitates oligomerisation of NPR1, whereas thioredoxins support reductive monomer release, a reaction that is boosted by SA (Figure 2). After identifying the crucial components, the next challenge will be to investigate how this intricate balance is achieved.

S-nitrosylation modulates phytohormonal signalling

NPR1 not only mediates defence gene expression, but also contributes to the suppression of JA-dependent responses in the cytosol in a so far unknown mode [32]. *S*-nitrosylation could thus contribute to the well-documented negative cross-talk between SA and JA signalling pathways. Moreover, allene oxide cyclase, a JA biosynthetic enzyme, has been shown to be *S*-nitrosylated during the hypersensitive response [26], which potentially represents another mechanism regulating oxylipin levels. In this connection, *S*-nitrosylation of salicylic acid binding protein 3 (SABP3) may also interfere with signal cross-talk by affecting lipid-derived signalling components as both carbonic anhydrase activity and SA-binding capacity of SABP3 are inhibited by *S*-nitrosylation [33]. Besides its apparent influence on SA/JA cross-talk, NO also affects ethylene, abscisic acid (ABA) and auxin signalling [34,35*,36]. *S*-nitrosylation of methionine adenosyltransferase 1 (MAT1; involved in ethylene biosynthesis) inhibits ethylene production [34], whereas NO is required for downstream responses to auxin and ABA [35*,36]. The interplay between different phytohormones and NO during innate immune reactions is an outstanding example of the emerging mutual influences and of how NO plays its regulatory role in vital functions of plants. ABA-dependent and SA-dependent stomatal closure upon pathogen recognition is mediated by NO; the virulence factor coronatine, a structural and functional mimic of JA, can reverse the effect and lead to stomatal reopening. These data implicate the involvement of at least three phytohormones (or mimics thereof) in pathogen-induced stomatal movements and suggest NO as a key mediator of these hormone responses [35*]. Although the role of NO in stomatal closure remains to be defined, *S*-nitrosylation of K⁺ outward rectifying chan-

nels supports a potential role of this posttranslational modification in modulating stomatal movements [37].

S-nitrosylation controls cell death

NO's control over Arabidopsis metacaspase 9, a potential executioner of programmed cell death [38*], is another intriguing example of how NO exerts its physiological functions through *S*-nitrosylation. Data indicate that this modification keeps the zymogen inactive under normal physiological conditions, inhibiting the autoproteolytic activity of prometacaspase 9. *S*-nitrosylation of Arabidopsis cytosolic glyceraldehyde 3-phosphate dehydrogenase (GAPDH) could also play a role in the regulation of cell death, as it causes GAPDH inactivation and potentially induces translocation to the nucleus [39]. In animal systems, *S*-nitrosylated GAPDH binds to and stabilises Siah1 (an E3 ubiquitin ligase), followed by translocation to the nucleus, where Siah1 degrades nuclear proteins. These events finally lead to apoptosis [40]. A similar scenario, though conceivable, still remains to be demonstrated in plant cells. *S*-nitrosylation seems to have opposing roles in controlling programmed cell death: on the one hand, *S*-nitrosylation promotes the execution of programmed cell death by altering GAPDH function and on the other hand, *S*-nitrosylation of prometacaspase 9 keeps this putative executioner of cell death inactive. However, the kinetics of these signalling reactions suggests the involvement of enzymatic mechanisms [41]. Hence, activation of metacaspase 9 under conditions of increased NO production may depend on enzymatic denitrosylation. In this context, thioredoxins clearly deserve attention in the future ([42*], cf. NPR1).

This overview of *S*-nitrosylation already gives an impression of the emerging complexity of NO-dependent signalling in plants. But there is more to come: Arabidopsis PrxIII has been found to be *S*-nitrosylated during the hypersensitive disease resistance response, resulting in not only inhibition of its hydroperoxide-reducing peroxidase activity, but also reduction of its ability to detoxify ONOO⁻ [22**]. This inhibition leads to increased ONOO⁻-dependent nitrotyrosine formation. Thus, *S*-nitrosylation can modulate a central point of convergence for ROS-dependent and NO-dependent signalling pathways in response to several stresses by deactivating a crucial component of the cellular antioxidant system.

Protein tyrosine nitration

Protein tyrosine nitration results from a chemical reaction, in which ONOO⁻ adds a nitro group in *ortho* position to the aromatic ring of tyrosine residues [43,44]. The added nitro group can alter protein function and conformation when placed on a relevant tyrosine residue and can increase the proteins' susceptibility to proteolysis [45,46].

In animals, nitration of tyrosine residues is often considered a marker for certain pathological conditions, but

has recently been suggested to be relevant to redox signalling [47*,48]. In plants, tyrosine nitration appears to be physiologically relevant during biotic and abiotic stresses [22**,49,50]. However, any physiological reaction must meet certain criteria to be considered a signalling process: (i) the signal compound is produced at strictly controlled rates; (ii) the induced reaction is reversible and/or the effects transient; (iii) the resulting modification (e.g. of a target protein) causes alterations of cell function(s). On the basis of these criteria, what arguments can be put forward to qualify tyrosine nitration as a signalling process in plants? (i) ONOO⁻ production is not under direct enzymatic control, but nonetheless seems to be tightly regulated (see 'Regulation of NO bioavailability'). The transient increase of ONOO⁻ levels in elicitor challenged tobacco cell culture supports this notion [50]. (ii) Recent studies report a transient increase of tyrosine-nitrated proteins during the disease resistance response [22**,50,51]. (iii) The only indications for functional modifications by tyrosine nitration come from the animal kingdom, where both enzyme inhibition and activation have been reported [52–54]. In plants, only one study provided an indirect proof of inhibition of glutathione *S*-transferase by RNS that correlated with increased nitrotyrosine levels [55]. We have recently conducted a proteomic study to identify tyrosine nitrated proteins in Arabidopsis plants challenged with an incompatible bacterial pathogen [51]. Interestingly, several of the proteins with increased nitration levels during the hypersensitive resistance response are also nitrated in unchallenged plants. Thus, tyrosine nitration could play a role under physiological as well as stress conditions. Independently, we found that NtMEK2, a tobacco MAPKK, can be nitrated *in vitro*, resulting in loss of its activity (E Vandelle *et al.*, unpublished). In animal systems, ONOO⁻ is known to interfere with phosphorylation cascades via nitration of relevant tyrosine residues in target proteins, mimicking or preventing tyrosine phosphorylation [47*]. Furthermore, several protein kinases, comprising all components of the MAPK module [47*], have been shown to be modulated by tyrosine nitration, being either activated [56,57] or inhibited [58,59]. On the basis of these findings, regulatory mechanisms controlling the MAPK module by tyrosine nitration are also plausible in plants, again pointing to RNS as crucial modulators of signal transduction.

Taken together, the above-mentioned data support our view that tyrosine nitration is a physiologically relevant posttranslational modification in plants.

Metal nitrosylation

NO can react with most transition metals, but its interactions with iron are probably the most relevant biologically [41,60]. The most prominent examples for modulation of enzymatic activity by NO in animals are associated with haem nitrosylation, such as inhibition of cytochrome *c* oxidase and NOS, regulation of diverse

cytochrome P450s and activation of soluble guanylate cyclase (sGC). sGC is the prototypic NO sensor and NO-modified protein in animals [61]. sGC is activated via haem nitrosylation and is responsible for the majority of physiological responses to NO. By contrast, virtually nothing is known about this NO-dependent modification in plants. The only exceptions are plant haemoglobins (cf. 'Regulation of NO bioavailability'). However, in this case the occurring reaction constitutes an NO scavenging mechanism and not a modification of enzymatic function. On the basis of chemical considerations, plant cytochrome P450s seem likely targets of metal nitrosylation. Given the functional diversity of plant cytochrome P450s, intriguing connections between NO-related signalling and different metabolic processes could emerge.

The missing link to cGMP

In animals, NO can initiate its biological effects through the activation of sGC and associated increase in the levels of the second messenger cGMP. Both a transient increase in cGMP and its involvement in several processes have also been demonstrated in plants [62]. Indeed, pharmacological and biochemical approaches showed that cGMP is involved in NO-dependent signalling, gene transcription modulation, root growth and gravitropism, pollen tube growth and orientation, hormone-dependent responses, stomatal opening and responses to biotic and abiotic stresses. Hence, the physiological role of this second messenger is unequivocally recognised. Although several potential GCs have been identified using bioinformatics tools, their relevance as cGMP sources still awaits demonstration. In fact, the overexpression of AtGC1 in *E. coli* [63] and activity analysis of the putative GC catalytic centre of the brassinosteroid receptor AtBRI1 *in vitro* [64] showed only low activity compared with the animal sGC [65], questioning their relevance as plant GCs. Supporting this observation, despite the capacity of these proteins to produce cGMP from GTP *in vitro*, our experiments did not reveal any increase in cGMP level in Arabidopsis plants overexpressing AtGC1 (J Hussain *et al.*, unpublished data). By contrast, we have produced Arabidopsis plants expressing the α -subunit and β -subunit of mammalian sGC that display an up to 40-fold increase in cGMP levels compared with wild-type plants (J Hussain *et al.*, unpublished). This has made a genetic tool available to study the impact of this second messenger in plants. The endogenous cGMP production systems, however, remain unclear.

Conclusions

We continue to gain new knowledge on NO-related signalling. However, the data available are still far from offering a comprehensive and consistent picture of NO function in plants. The lack of genetic tools substantially hinders research on NO production and functions. New findings, however, underline the importance of NO in plant cell physiology and the complexity of NO-related

signalling networks. Particularly, the impact of *S*-nitrosylation on protein function has been clearly demonstrated, plus different modes of NO-mediated signalling are gaining increasing attention. On the basis of the current status of research we can expect the functional relevance of other NO-mediated posttranslational modifications and/or second messengers to be demonstrated soon.

Acknowledgement

This work was supported by a grant to M.D. from the Ministero dell'Università e della Ricerca in the framework of the program 'Components of the nitric oxide signalling pathways in plants'.

References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
 - of outstanding interest
1. Delledonne M, Xia Y, Dixon RA: **Lamb C: nitric oxide functions as a signal in plant disease resistance.** *Nature* 1998, **394**:585-588.
 2. Wilson ID, Neill SJ, Hancock JT: **Nitric oxide synthesis and signalling in plants.** *Plant Cell Environ* 2008, **31**:622-631.
This review, together with [4*], gives a more detailed overview of NO production in plants and some aspects of signalling that are not covered here due to space limitations.
 3. Nathan C: **The moving frontier in nitric oxide-dependent signalling.** *Sci STKE* 2004, **2004**:pe52.
 4. Besson-Bard A, Pugin A, Wendehenne D: **New insights into nitric oxide signaling in plants.** *Annu Rev Plant Biol* 2008, **59**:21-39.
As [2*], this review describes NO signalling and production in plants in more detail; both references also contain extensive bibliographies.
 5. Moreau M, Lee GI, Wang Y, Crane BR, Klessig DF: **AtNOS/AtNOA1 is a functional *Arabidopsis thaliana* cGTPase and not a nitric-oxide synthase.** *J Biol Chem* 2008, **283**:32957-32967.
After a long controversy, this study finally demonstrated that AtNOS1/AtNOA1 is not a nitric oxide synthase, but a chloroplastic GTPase involved in proper ribosome assembly.
 6. Flores-Pérez U, Sauret-Güeto S, Gas E, Jarvis P, Rodríguez-Concepción M: **A mutant impaired in the production of plastome-encoded proteins uncovers a mechanism for the homeostasis of isoprenoid biosynthetic enzymes in *Arabidopsis* plastids.** *Plant Cell* 2008, **20**:1303-1315.
Together with [4*], this study ended the controversy about AtNOS1/AtNOA1/AtRif1. The gene was identified independently in a screening for fosmidomycin resistance. Levels of methylerythritol pathway enzymes are upregulated in the mutant.
 7. Bright J, Desikan R, Hancock JT, Weir IS, Neill SJ: **ABA-induced NO generation and stomatal closure in *Arabidopsis* are dependent on H₂O₂ synthesis.** *Plant J* 2006, **45**:113-122.
 8. Modolo LV, Augusto O, Almeida IMG, Pinto-Maglio CAF, Oliveira HC, Seligman K, Salgado I: **Decreased arginine and nitrite levels in nitrate reductase-deficient *Arabidopsis thaliana* plants impair nitric oxide synthesis and the hypersensitive response to *Pseudomonas syringae*.** *Plant Sci* 2006, **171**:34-40.
 9. Flores T, Todd CD, Tovar-Mendez A, Dhanoa PK, Correa-Aragunde N, Hoyos ME, Brownfield DM, Mullen RT, Lamattina L, Polacco JC: **Arginase-negative mutants of *Arabidopsis* exhibit increased nitric oxide signaling in root development.** *Plant Physiol* 2008, **147**:1936-1946.
 10. He Y, Tang R-H, Hao Y, Stevens RD, Cook CW, Ahn SM, Jing L, Yang Z, Chen L, Guo F *et al.*: **Nitric oxide represses the *Arabidopsis* floral transition.** *Science* 2004, **305**:1968-1971.
 11. Lee U, Wie C, Fernandez BO, Feelisch M, Vierling E: **Modulation of nitrosative stress by *S*-nitrosoglutathione reductase is critical for thermotolerance and plant growth in *Arabidopsis*.** *Plant Cell* 2008, **20**:786-802.
 12. Morot-Gaudry-Talarmain Y, Rockel P, Moureaux T, Quilleré I, Leydecker MT, Kaiser WM, Morot-Gaudry JF: **Nitrite accumulation and nitric oxide emission in relation to cellular signaling in nitrite reductase antisense tobacco.** *Planta* 2002, **215**:708-715.
 13. Ahlfors R, Brosché M, Kollist H, Kangasjärvi J: **Nitric oxide modulates ozone-induced cell death, hormone biosynthesis and gene expression in *Arabidopsis thaliana*.** *Plant J* 2009, **58**:1-12.
 14. Avila EL, Brown M, Pan S, Desikan R, Neill SJ, Girke T, Surpin M, Raikhel NV: **Expression analysis of *Arabidopsis* vacuolar sorting receptor 3 reveals a putative function in guard cells.** *J Exp Bot* 2008, **59**:1149-1161.
 15. Iwakiri Y, Satoh A, Chatterjee S, Toomre DK, Chalouni CM, Fulton D, Groszmann RJ, Shah VH, Sessa WC: **Nitric oxide synthase generates nitric oxide locally to regulate compartmentalized protein *S*-nitrosylation and protein trafficking.** *Proc Natl Acad Sci USA* 2006, **103**:19777-19782.
 16. Harris LK, McCormick J, Cartwright JE, Whitley GSJ, Dash PR: ***S*-nitrosylation of proteins at the leading edge of migrating trophoblasts by inducible nitric oxide synthase promotes trophoblast invasion.** *Exp Cell Res* 2008, **314**:1765-1776.
 17. Perazolli M, Dominici P, Romero-Puertas MC, Zago E, Zeier J, Sonoda M, Lamb C, Delledonne M: ***Arabidopsis* nonsymbiotic hemoglobin AHb1 modulates nitric oxide bioactivity.** *Plant Cell* 2004, **16**:2785-2794.
 18. Zeier J, Delledonne M, Mishina T, Severi E, Sonoda M, Lamb C: **Genetic elucidation of nitric oxide signaling in incompatible plant-pathogen interactions.** *Plant Physiol* 2004, **136**:2875-2886.
 19. Liu L, Hausladen A, Zeng M, Que L, Heitman J, Stamler JS: **A metabolic enzyme for *S*-nitrosothiol conserved from bacteria to humans.** *Nature* 2001, **410**:490-494.
 20. Feechan A, Kwon E, Yun B-W, Wang Y, Pallas JA, Loake GJ: **A central role for *S*-nitrosothiols in plant disease resistance.** *Proc Natl Acad Sci USA* 2005, **102**:8054-8059.
 21. Rustérucci C, Espunya MC, Díaz M, Chabannes M, Martínez MC: ***S*-nitrosoglutathione reductase affords protection against pathogens in *Arabidopsis*, both locally and systemically.** *Plant Physiol* 2007, **143**:1282-1292.
 22. Romero-Puertas MC, Laxa M, Matté A, Zaninotto F, Finkemeier I, Jones AME, Perazolli M, Vandelle E, Dietz K-J, Delledonne M: ***S*-nitrosylation of peroxiredoxin II E promotes peroxynitrite-mediated tyrosine nitration.** *Plant Cell* 2007, **19**:4120-4130.
Peroxiredoxin II E is involved in peroxynitrite detoxification. This activity is inhibited by *S*-nitrosylation of the enzyme, which leads to increased protein nitration by peroxynitrite. The combined functions of hydroperoxide and peroxynitrite detoxification suggest that PrxII E modulates both ROS and NO signalling.
 23. Delledonne M, Zeier J, Marocco A, Lamb C: **Signal interactions between nitric oxide and reactive oxygen intermediates in the plant hypersensitive disease resistance response.** *Proc Natl Acad Sci USA* 2001, **98**:13454-13459.
 24. Stamler JS, Lamas S, Fang FC: **Nitrosylation: the prototypic redox-based signaling mechanism.** *Cell* 2001, **106**:675-683.
 25. Lindermayr C, Saalbach G, Durner J: **Proteomic identification of *S*-nitrosylated proteins in *Arabidopsis*.** *Plant Physiol* 2005, **137**:921-930.
 26. Romero-Puertas MC, Campostrini N, Matté A, Righetti PG, Perazolli M, Zolla L, Roepstorff P, Delledonne M: **Proteomic analysis of *S*-nitrosylated proteins in *Arabidopsis thaliana* undergoing hypersensitive response.** *Proteomics* 2008, **8**:1459-1469.
 27. Ferrarini A, De Stefano M, Baudouin E, Pucciariello C, Polverari A, Puppo A, Delledonne M: **Expression of *Medicago truncatula* genes responsive to nitric oxide in pathogenic and symbiotic conditions.** *Mol Plant Microbe Interact* 2008, **21**:781-790.

28. Grün S, Lindermayr C, Sell S, Durner J: **Nitric oxide and gene regulation in plants.** *J Exp Bot* 2006, **57**:507-516.
29. Palmieri MC, Sell S, Huang X, Scherf M, Werner T, Durner J, Lindermayr C: **Nitric oxide-responsive genes and promoters in *Arabidopsis thaliana*: a bioinformatics approach.** *J Exp Bot* 2008, **59**:177-186.
30. Serpa V, Vernal J, Lamattina L, Grotewold E, Cassia R, Terenzi H: **Inhibition of AtMYB2 DNA-binding by nitric oxide involves cysteine S-nitrosylation.** *Biochem Biophys Res Commun* 2007, **361**:1048-1053.
31. Tada Y, Spoel SH, Pajerowska-Mukhtar K, Mou Z, Song J, Wang C, Zuo J, Dong X: **Plant immunity requires conformational charges of NPR1 via S-nitrosylation and thioredoxins.** *Science* 2008, **321**:952-956.
- S-nitrosylation is involved the control of NPR1 oligomerisation and localisation and thus influences the activation of SA-dependent defence responses.
32. Spoel SH, Koornneef A, Claessens SMC, Korzelius JP, Van Pelt JA, Mueller MJ, Buchala AJ, Métraux J-P, Brown R, Kazan K *et al.*: **NPR1 modulates cross-talk between salicylate- and jasmonate-dependent defense pathways through a novel function in the cytosol.** *Plant Cell* 2003, **15**:760-770.
33. Wang Y-Q, Feechan A, Yun B-W, Shafiei R, Hofmann A, Taylor P, Xue P, Yang F-Q, Xie Z-S, Pallas JA *et al.*: **S-nitrosylation of AtSABP3 antagonizes the expression of plant immunity.** *J Biol Chem* 2009, **284**:2131-2137.
34. Lindermayr C, Saalbach G, Bahnweg G, Durner J: **Differential inhibition of *Arabidopsis* methionine adenosyltransferases by protein S-nitrosylation.** *J Biol Chem* 2006, **281**:4285-4291.
35. Melotto M, Underwood W, Koczan J, Nomura K, He SY: **Plant stomata function in innate immunity against bacterial invasion.** *Cell* 2006, **126**:969-980.
- The authors show that NO is involved in SA-dependent and ABA-dependent stomatal closure upon pathogen recognition. The virulence factor coronatine is involved in reopening of stomata and thus circumvention of this mechanism of innate immunity. This study gives an impressive example of the intricate interplay of different signals.
36. Correa-Aragunde N, Graziano M, Lamattina L: **Nitric oxide plays a central role in determining lateral root development in tomato.** *Planta* 2004, **218**:900-905.
37. Sokolovskii S, Blatt MR: **Nitric oxide block of outward-rectifying K⁺ channels indicates direct control by protein nitrosylation in guard cells.** *Plant Physiol* 2004, **136**:4275-4284.
38. Belenghi B, Romero-Puertas MC, Vercammen D, Brackenhier A, Inzé D, Delledonne M, Van Breusegem F: **Metacaspase activity of *Arabidopsis thaliana* is regulated by S-nitrosylation of a critical cysteine residue.** *J Biol Chem* 2007, **282**:1352-1358.
- The authors provide evidence for the inhibition of autoproteolytic activity of prometacaspase 9 by S-nitrosylation, whereas the mature protease is not inhibited by NO. Thus, steady-state S-nitrosylation in unchallenged plants may be a mechanism to keep executioners of programmed cell death inactive.
39. Holtgreve S, Gohlke J, Starmann J, Druce S, Klocke S, Altmann B, Wojtera J, Lindermayr C, Scheibe R: **Regulation of plant cytosolic glyceraldehyde 3-phosphate dehydrogenase isoforms by thiol modifications.** *Physiol Plant* 2008, **133**:211-228.
40. Hara MR, Agrawal N, Kim SF, Cascio MB, Fujimuro M, Ozeki Y, Takahashi M, Cheah JH, Tankou SK, Hester LD *et al.*: **S-nitrosylated GAPDH initiates apoptotic cell death by nuclear translocation following Siah1 binding.** *Nat Cell Biol* 2005, **7**:665-674.
41. Forman HJ, Fukuto JM, Miller T, Zhang H, Rinna A, Levy S: **The chemistry of cell signaling by reactive oxygen and nitrogen species and 4-hydroxynonenal.** *Arch Biochem Biophys* 2008, **477**:183-195.
42. Benhar M, Forrester MT, Hess DT, Stamler JS: **Regulated protein denitrosylation by cytosolic and mitochondrial thioredoxins.** *Science* 2008, **320**:1050-1054.
- The authors demonstrated that thioredoxins act as denitrosylating enzymes in animal systems. Besides their role in protection against oxidative damage, thioredoxins potentially contribute to NO-related signal transduction by modulating basal and stimulus induced denitrosylation.
43. Radi R: **Nitric oxide, oxidants, and protein tyrosine nitration.** *Proc Natl Acad Sci USA* 2004, **101**:4003-4008.
44. Rubbo H, Radi R: **Protein and lipid nitration: role in redox signaling and injury.** *Biochim Biophys Acta* 2008, **1780**:1318-1324.
45. Grune T, Blasig IE, Sitte N, Roloff B, Haseloff R, Davies KJA: **Peroxynitrite increases the degradation of aconitase and other cellular proteins by proteasome.** *J Biol Chem* 1998, **273**:10857-10862.
46. Souza JM, Choi I, Chen Q, Weisse M, Daikhin E, Yudkoff M, Obin M, Ara J, Horwitz J, Ischiropoulos H: **Proteolytic degradation of tyrosine nitrated proteins.** *Arch Biochem Biophys* 2000, **380**:360-366.
47. Monteiro HP, Arai RJ, Travassos LR: **Protein tyrosine phosphorylation and protein tyrosine nitration in redox signaling.** *Antioxid Redox Signal* 2008, **10**:843-890.
- This review presents a very good survey of the competition/cooperation between tyrosine phosphorylation and tyrosine nitration in animals. Emphasis is placed on the role of nitration in modulating phosphorylation-mediated cellular events by mimicking or inhibiting tyrosine-phosphorylation.
48. Aulak KS, Koeck T, Crabb JW, Stuehr DJ: **Proteomic method for identification of tyrosine-nitrated proteins.** *Methods Mol Biol* 2004, **279**:151-165.
49. Corpas FJ, Chaki M, Fernandez-Ocana A, Valderrama R, Palma JM, Carreras A, Begara-Morales JC, Airaki M, del Rio LA, Barroso JB: **Metabolism of reactive nitrogen species in pea plants under abiotic stress conditions.** *Plant Cell Physiol* 2008, **49**:1711-1722.
50. Saito S, Yamamoto-Katou A, Yoshioka H, Doke N, Kawakita K: **Peroxynitrite generation and tyrosine nitration in defense responses in tobacco BY-2 cells.** *Plant Cell Physiol* 2006, **47**:689-697.
51. Cecconi D, Orzetti S, Vandelle E, Rinalducci S, Zolla L, Delledonne M: **Protein nitration during defence response in *Arabidopsis thaliana*.** *Electrophoresis* 2009, doi:10.1002/elps.200800826.
52. Neumann H, Hazen JL, Weinstein J, Mehl RA, Chin JW: **Genetically encoding protein oxidative damage.** *J Am Chem Soc* 2008, **130**:4028-4033.
53. Wong PSY, Eiserich JP, Reddy S, Lopez CL, Cross CE, van der Vliet A: **Inactivation of glutathione S-transferases by nitric oxide-derived oxidants: exploring a role for tyrosine nitration.** *Arch Biochem Biophys* 2001, **394**:216-228.
54. Cassina AM, Hodara R, Souza JM, Thomson L, Castro L, Ischiropoulos H, Freeman BA, Radi R: **Cytochrome c nitration by peroxynitrite.** *J Biol Chem* 2000, **275**:21409-21415.
55. Gong Y-W, Yuan Y-J: **Nitric oxide mediates inactivation of glutathione S-transferase in suspension culture of *Taxus cuspidata* during shear stress.** *J Biotechnol* 2006, **123**:185-192.
56. Zhang P, Wang Y-Z, Kagan E, Bonner JC: **Peroxynitrite targets the epidermal growth factor receptor, Raf-1, and MEK independently to activate MAPK.** *J Biol Chem* 2000, **275**:22479-22486.
57. Pinzar E, Wang T, Garrido MR, Xu W, Levy P, Bottari SP: **Angiotensin II induces tyrosine nitration and activation of ERK1/2 in vascular smooth muscle cells.** *FEBS Lett* 2005, **579**:5100-5104.
58. Schieke S, Briviba K, Klotz L-O, Sies H: **Activation pattern of mitogen-activated protein kinases elicited by peroxynitrite: attenuation by selenite supplementation.** *FEBS Lett* 1999, **448**:301-303.
59. Kinjo N, Kawanaka H, Akahoshi T, Yamaguchi S, Yoshida D, Aneagawa G, Konishi K, Tomikawa M, Tanoue K, Tarnawski A *et al.*: **Significance of ERK nitration in portal hypertensive gastropathy and its therapeutic implications.** *Am J Physiol Gastrointest Liver Physiol* 2008, **295**:G1016-1024.
60. Cooper CE: **Nitric oxide and iron proteins.** *Biochim Biophys Acta* 1999, **1411**:290-309.
61. Poulos TL: **Soluble guanylate cyclase.** *Curr Opin Struct Biol* 2006, **16**:736-743.

62. Meier S, Gehring C: **Emerging roles in plant biotechnology for the second messenger cGMP—guanosine 3', 5'-cyclic monophosphate.** *Afr J Biotechnol* 2006, **5**:1687-1692.
63. Ludidi N, Gehring C: **Identification of a novel protein with guanylyl cyclase activity in *Arabidopsis thaliana*.** *J Biol Chem* 2003, **278**:6490-6494.
64. Kwezi L, Meier S, Mungur L, Ruzvidzo O, Irving H, Gehring C: **The *Arabidopsis thaliana* brassinosteroid receptor (AtBRI1) contains a domain that functions as a guanylyl cyclase *in vitro*.** *PLoS ONE* 2007, **2**:e449.
65. Winger JA, Marletta MA: **Expression and characterization of the catalytic domains of soluble guanylate cyclase: interaction with the heme domain.** *Biochemistry* 2005, **44**:4083-4090.